

FROM TOPOLOGICAL QUANTUM MECHANICS TO THE PHYSICS OF SPIN-ONE PARTICLES

Romain Attal and Laurent Baulieu *

LPTHE Universités Pierre et Marie Curie et Denis Diderot

Laboratoire associé au CNRS, URA 280

4 place Jussieu, Tour 16, 1er étage

F-75252 Paris Cedex 05, France.

Abstract: We give an example of topological theory whose Hilbert space contains physical objects: the $N = 2$ supersymmetric Lagrangian for spin-one particles moving in D -dimensional space-time is a topological model in a space with two more dimensions. The equivalence is still valid in a curved space-time. As an application, we calculate the deviation of spin-one particles in a Schwarzschild background.

*e-mail address: attal@lpthe.jussieu.fr baulieu@lpthe.jussieu.fr

1 Introduction.

Supersymmetric quantum mechanics has specific applications: it can be used to describe the dynamics of spinning point particles [1] as well as to compute a certain number of topological invariants of the target space [2]. Moreover it gives an insight into the structure of superstring theory, and has opened the way to its modern formulation [3] [4].

In this paper we point out an example showing that topological quantum theories may exhibit a phase with a Hilbert space made of particle degrees of freedom. We implement the idea that the fundamental symmetries of nature could be originally of the topological type and that the observed gauge symmetries would be obtained by gauge-fixing the huge topological symmetry in a BRST invariant way, leaving therefore an $N = 2$ supersymmetric theory.

Our example is the description of spin-one particles by a $N = 2$ supersymmetric action. The use of anticommuting variables to describe spinning particles was introduced in [1]. Then, it was found that local supersymmetry of rank $2S$ on the worldline is necessary to describe consistently a particle of spin S . The resulting constrained system [5] [6] requires a careful gauge-fixing of the einbein and the gravitini. One obtains eventually a tractable Lagrangian formulation [7], [8]. (There are many references on the subject, of which we quote very few).

In this work we interpret local supersymmetry on the worldline as a residue of a more fundamental topological symmetry, defined in a target-space with two extra dimensions. The word “topological” has to be understood in the sense that one does the BRST invariant gauge-fixing of a classical action, such as a characteristic number of the base manifold, which is invariant under arbitrary infinitesimal deformations of the fields [9]. One of the coordinates is eventually identified as the einbein on the worldline. Other fields must be introduced to enforce the topological BRST invariance. They can be eliminated by their equations of motion and decouple from the physical sector. To obtain in a natural way a nowhere vanishing einbein, we use a disconnected higher dimensional target-space

where the hyperplane $\{e = 0\}$ is a priori extracted. Thus, one introduces some topology before any gauge-fixing. Two disconnected topological sectors exist, $\{e > 0\}$ and $\{e < 0\}$, which correspond to the prescription $\pm i\epsilon$ for the propagators. It is fundamental that the gauge functions be compatible with the topology of space: they must induce a potential which rejects the trajectories from the hyperplane $\{e = 0\}$.

The paper is organized as follows. We first review the supersymmetric description of a relativistic spinning particle in a Riemannian space-time. Then we consider the case of $N = 2$ supersymmetry and show our main result: the link between the supersymmetric description of scalar or spin-one particles and topological quantum mechanics in a higher dimensional target-space. In the last section, we verify that the constraints of the theory identify its physical content and illustrate the result by computing the deviation of the trajectories from geodesics due to the interactions between geometry and spin.

2 Worldline supergravity for a spinning particle and its gauge fixing.

Consider a spin- S particle in a D -dimensional space-time. Classically, it follows a world-line whose coordinates $X^\mu(\tau)$ are parametrized by a real number τ . If the particle is massive, a natural choice of this parameter is the proper-time. The idea originating from [1] is to describe the spin of the particle by assigning to each value of τ a vector with anticommuting coordinates $\Psi_i^\mu(\tau)$ where the vector index μ runs between 1 and D and i between 1 and $2S$. Indeed, in the case of a flat space-time and spin one-half, the Lagrangian density introduced in [1] is

$$\mathcal{L} = \frac{1}{2}(\dot{X}^2(\tau) - \Psi^\mu(\tau)\dot{\Psi}_\mu(\tau)) \quad (2.1)$$

where the dot $\dot{}$ means ∂_τ , τ being a parametrization of the worldline. Upon canonical quantization $\Psi^\mu(\tau)$ is replaced by a τ -independent operator $\hat{\Psi}^\mu$ which satisfies anticommutation relations

$$\{\hat{\Psi}^\mu, \hat{\Psi}_\nu\}_+ = 2\delta_\nu^\mu \quad (2.2)$$

The Hamiltonian is

$$H = \frac{1}{2}p^2 = \frac{1}{2}Q^2 \quad (2.3)$$

with $Q = p_\mu \hat{\Psi}^\mu$. Due to (2.2) the $\hat{\Psi}$'s can be represented by Dirac matrices and Q is the free Dirac operator. Q commutes with H and it makes sense to consider the restriction of the Hilbert space to the set of states $|\varphi\rangle$ satisfying

$$Q|\varphi\rangle = 0 \quad (2.4)$$

By definition of Q , this equation means that the $|\varphi\rangle$ are the states of a massless spin one-half particle. The extension to the case of a massive particle implies the introduction of an additional Grassmannian variable Ψ^{D+1} and the generalization of $\mathcal{L}$ to

$$\mathcal{L} = \frac{1}{2}(\dot{X}^2(\tau) - \Psi^\mu(\tau)\dot{\Psi}_\mu(\tau) - \Psi^{D+1}(\tau)\dot{\Psi}^{D+1}(\tau) + m^2) \quad (2.5)$$

(Formally, $\dot{X}^{D+1} \rightarrow m$), so that

$$H = \frac{1}{2}(p^2 - m^2) = \frac{1}{2}Q^2 \quad (2.6)$$

with

$$Q = p_\mu \hat{\Psi}^\mu + m\hat{\Psi}^{D+1} \quad (2.7)$$

and one has in addition to (2.2)

$$\{\hat{\Psi}^{D+1}, \hat{\Psi}^{D+1}\}_+ = -2 \quad \{\hat{\Psi}^\mu, \hat{\Psi}^{D+1}\}_+ = 0 \quad (2.8)$$

The condition (2.4) is now the free Dirac equation for a spin one-half particle of mass m , multiplied by $\hat{\Psi}^{D+1}$. The generalization to the case of an arbitrary spin is obtained by duplicating $2S$ times the components of Ψ , $\Psi^\mu \rightarrow \Psi_i^\mu, 1 \leq i \leq 2S$, as can be seen by constructing the representations of $SO(D)$ by suitable tensor products of spin one-half representations [12] [13].

To understand the constraint (2.4), it is in fact necessary to promote the global supersymmetry of the action, corresponding to the commutation of H and Q , into a local supersymmetry. Indeed, when time flows, the state of the particle must evolve

from a solution of the Dirac equation to another solution of this equation, without any possibility to collapse in an unphysical state (out of $Ker(Q)$). A natural way to reach such a unitarity requirement is to impose the supersymmetry independently for all values of τ , that is, to gauge the supersymmetry on the worldline. In this way, the condition (2.4) appears as the definition of physical states in a gauge theory with generator Q which ensures unitarity, like the transversality condition of gauge bosons in ordinary Yang-Mills theory. For consistency, the diffeomorphism invariance on the worldline must be also imposed since the commutator of two supersymmetry transformations contains a diffeomorphism. One thus introduces gauge fields for these symmetries, the einbein $e(\tau)$ and the (anticommuting) gravitino $\alpha(\tau)$. By minimal coupling on the worldline, (2.5) is thus generalized to the following Lagrangian which is locally supersymmetric and reparametrization invariant, up to a pure derivative with respect to τ :

$$\mathcal{L} = \frac{1}{2} \left(e^{-1} \dot{X}^2 - \Psi(\dot{\Psi} + \alpha e^{-1} \dot{X}) - \Psi^{D+1}(\dot{\Psi}^{D+1} + m\alpha) + em^2 \right) \quad (2.9)$$

(In this section we omit the vector and spin indices). Formally, $\dot{X}^{D+1} \rightarrow me$. The transformation laws of e and α are those of one-dimensional supergravity of rank $2S$.

The gauge-fixing $e(\tau) = 1$ and $\alpha(\tau) = 0$ identifies (2.5) and (2.9), up to Faddeev-Popov ghost terms. These ghost terms have a supersymmetric form $b\dot{c} + \beta\dot{\gamma}$. They decouple effectively, since their effect is to multiply all the amplitudes by a ratio of determinants, independent of the metric in space-time. This gauge-fixing is however inconsistent because it is too strong, since the Lagrangian is gauge invariant only up to boundary terms. Therefore, given a general gauge transformation, one must put restrictions on its parameters to get the invariance of the action, and there are not enough degrees of freedom in the symmetry to enforce the gauge $e(\tau) = 1$ and $\alpha(\tau) = 0$. One can at most set $e(\tau) = e_0$ and $\alpha(\tau) = \alpha_0$, letting the constants $e_0 > 0$ and α_0 free, that is, doing an ordinary integration over e_0 and α_0 in the path integral after the gauge-fixing [7]. This yields the following partition function for the theory

$$Z = \int_0^\infty de_0 \int d\alpha_0 \int [dX(\tau)][d\Psi(\tau)] \exp - \int_0^1 d\tau \mathcal{L}_0 \quad (2.10)$$

with

$$\mathcal{L}_0 = \frac{1}{2} \left(e_0^{-1} \dot{X}^2 + e_0 m^2 - \Psi(\dot{\Psi} + \alpha_0 e_0^{-1} \dot{X}) - \Psi^{D+1}(\dot{\Psi}^{D+1} + m\alpha_0) \right) \quad (2.11)$$

Using the Lagrangian (2.5) instead of (2.11) implies that one misses crucial spin-orbit interactions described by the Grassmannian integration over the constant α_0 which induces the fermionic constraint $\int d\tau (\Psi \dot{X} + m e_0 \Psi^{D+1}) = 0$. The use of (2.5) leads indeed to a spin-zero particle propagator while (2.11) leads to the expected spin one-half propagator. One gets the $\pm i\epsilon$ propagators depending on the choice of the integration domain $\{e_0 > 0\}$ or $\{e_0 < 0\}$. Notice that the e -dependence of the Lagrangian (2.5) gives a negligible weight in the path integral (2.10) to the trajectories with points near the hyperplane $\{e_0 = 0\}$. The integration over e_0 and α_0 has a simple interpretation in Hamiltonian formalism. The Hamiltonian associated to (2.11) is

$$\begin{aligned} H &= \frac{e_0}{2}(p^2 - m^2) + \frac{\alpha_0}{2}(p_\mu \hat{\Psi}^\mu + m \hat{\Psi}^{D+1}) \\ &= \frac{e_0}{2}(p^2 - m^2) + \frac{\alpha_0}{2}Q \end{aligned} \quad (2.12)$$

The constants e_0 and α_0 are thus Lagrange multipliers which force the particle to satisfy the Klein-Gordon equation and the Dirac equation (or its higher spin generalizations $Q_i|\varphi\rangle = 0$). Observe that in Lagrangian formalism, the Klein-Gordon equation is not a consequence of the Dirac equation, due to the anticommutativity of Grassmann variables, and the two constraints $Q|\varphi\rangle = 0$ and $H|\varphi\rangle = 0$ must be used separately. Therefore, we have a theory where the Hamiltonian is a sum of constraints, which leads to known technical difficulties [5][6]. In Lagrangian formalism, supergravity on the worldline and its correct gauge-fixing take care of all details [7].

The above description is valid for a flat space-time. It can be generalized to the case where the particle moves in a curved space-time and/or couples to an external electromagnetic field, by minimal coupling in the target-space. The compatibility between the worldline diffeomorphism invariance and local supersymmetry with reparametrization invariance in the target-space for a general metric $g_{\mu\nu}$ is however possible only for $N \leq 2$ [12]. This phenomenon is possibly related to the limited number of consistent supergravities [14].

We now consider the case $N = 2$ and establish our main result in this paper: the link of the theory with a topological model.

3 The $N=2$ supersymmetric action as a topological action.

The $N = 2$ supersymmetric Lagrangian with a general background metric $g_{\mu\nu}$ is

$$\begin{aligned}\mathcal{L}_{SUSY} &= \frac{1}{2e}g_{\mu\nu}\dot{X}^\mu\dot{X}^\nu - \overline{\Psi}^\mu(g_{\mu\nu}\dot{\Psi}^\nu + e\Gamma_{\mu\nu\rho}\dot{X}^\nu\Psi^\rho) + e^{-1}g_{\mu\nu}\dot{X}^\mu(\overline{\Psi}^\nu\alpha + \overline{\alpha}\Psi^\nu) \\ &+ \frac{em^2}{2} - \overline{\Psi}^{D+1}\dot{\Psi}^{D+1} + m(\overline{\Psi}^{D+1}\alpha + \overline{\alpha}\Psi^{D+1}) \\ &- e^{-1}\overline{\alpha}\alpha\overline{\Psi}\Psi + \frac{e}{2}R_{\mu\nu\rho\sigma}\overline{\Psi}^\mu\Psi^\nu\overline{\Psi}^\rho\Psi^\sigma\end{aligned}\tag{3.1}$$

where Ψ and $\overline{\Psi}$ are independent Grassmannian coordinates. (Compare with [12]). The Lagrangian (3.1) has two local supersymmetries, with generators Q and $\overline{Q}$. An $O(2)$ symmetry between Ψ and $\overline{\Psi}$ can be enforced by introducing a single gauge field $f(\tau)$ and adding a term $f\overline{\Psi}\Psi$. However, no new information is provided, since one increases the symmetry by one generator, which is compensated by the introduction of the additional degree of freedom carried by f . The latter can indeed be gauge-fixed to zero and one recovers (3.1). Moreover, in view of identifying Ψ and $\overline{\Psi}$ as ghosts and antighosts, one wishes to freeze the symmetry between these two fields. We thus ignore the possibility of gauging the $O(2)$ symmetry. We will check in the next section that the Hilbert space associated to the Lagrangian (3.1) contains spin-one particles.

The Lagrangian (3.1) can be conveniently rewritten in first order formalism by introducing a Lagrange multiplier $b^\mu(\tau)$. One gets the equivalent form

$$\begin{aligned}\mathcal{L}_{SUSY} &\sim -\frac{e}{2}(g_{\mu\nu}b^\mu b^\nu - m^2) + g_{\mu\nu}b^\mu(\dot{X}^\nu + e\Gamma_{\rho\sigma}^\nu\overline{\Psi}^\rho\Psi^\sigma + \overline{\Psi}^\nu\alpha + \overline{\alpha}\Psi^\nu) \\ &- \overline{\Psi}^\mu(g_{\mu\nu}\dot{\Psi}^\nu + e\partial_\rho g_{\mu\nu}\dot{X}^\nu\Psi^\rho) - \Gamma_{\nu\rho\sigma}\overline{\Psi}^\rho\Psi^\sigma(\overline{\Psi}^\nu\alpha + \overline{\alpha}\Psi^\nu) \\ &- \overline{\Psi}^{D+1}\dot{\Psi}^{D+1} + m(\overline{\Psi}^{D+1}\alpha + \overline{\alpha}\Psi^{D+1}) - \frac{e}{2}\partial_\nu\Gamma_{\mu\rho\sigma}\overline{\Psi}^\mu\Psi^\nu\overline{\Psi}^\rho\Psi^\sigma\end{aligned}\tag{3.2}$$

(The symbol $\sim$ means that the two Lagrangians differ by a term which can be eliminated using an algebraic equation of motion, and, consequently, define the same quantum theory). For $e = 1$, $\alpha = \bar{\alpha} = 0$ and $\Psi^{D+1} = \bar{\Psi}^{D+1} = 0$, the Lagrangian (3.2) can be interpreted as the gauge-fixing of zero or of a term invariant under isotopies of the curve X [10]. In this interpretation the Ψ are topological ghosts and the $\bar{\Psi}$ are antighosts. The BRST graded differential operator s of the topological symmetry is defined by

$$\begin{aligned} sX^\mu &= \Psi^\mu \\ s\Psi^\mu &= 0 \\ s\bar{\Psi}^\mu &= b^\mu \\ sb^\mu &= 0 \end{aligned} \tag{3.3}$$

and the gauge-fixing Lagrangian is s -exact modulo a pure derivative

$$\mathcal{L}_{GF} = s(\bar{\Psi}_\mu(-\frac{1}{2}b^\mu + \dot{X}^\mu + \frac{1}{2}\Gamma^\mu_{\rho\sigma}\bar{\Psi}^\rho\Psi^\sigma)) \tag{3.4}$$

(Since $s^2 = 0$, $\mathcal{L}_{GF}$ is s -invariant.) To identify (3.1) as a topological Lagrangian, we must introduce new ingredients. We will enlarge the target-space with two additional components, and add a ghost of ghost. We will eventually identify one of the extra coordinates with the einbein e and the other one will be forced to vary in a Gaussian way around an arbitrary scale, with an arbitrary width. The gravitini α and $\bar{\alpha}$ of the effective worldline supergravity will be interpreted as ghosts of the topological symmetry. The $O(2)$ invariance corresponds to the ghost number conservation.

We consider a $(D+2)$ -dimensional space-time with coordinates $X^A = (X^\mu, X^{D+1} = e, X^{D+2})$. We exclude from the space the hyperplane $\{X^{D+1} = 0\}$ which yields two separated half-spaces, characterized by the value of $sign(e)$. We wish to define a partition function through a path integration over the curves $X^A(\tau)$, with a topological action which is invariant under the BRST symmetry associated to isotopies of this curve in each half-space. In other words we wish to construct an action by consistently gauge-fixing the topological Lagrangian $sign(e)$. In a way which is analogous to the case of topological Yang-Mills symmetry, where one gauge-fixes the second Chern class $\int Tr F^2$

[11], we combine the pure topological symmetry, with topological ghosts $\Psi_{top}^A(\tau)$, to the diffeomorphism symmetry on the curve, with Faddeev-Popov ghost $c(\tau)$. The apparent redundancy in the number of ghost variables $\Psi_{top}^A(\tau)$ and $c(\tau)$, which exceeds the number of bosonic classical variables, is counterbalanced by the introduction of a ghost of ghosts $\Phi(\tau)$ with ghost number two. The action of the BRST differential s is defined by

$$\begin{aligned}
sX^\mu &= \Psi_{top}^\mu + c\dot{X}^\mu = \Psi^\mu \\
se &= \Psi_{top}^e + c\dot{e} = 2\eta = \alpha + \dot{\Psi}^{D+1} \\
sX^{D+2} &= \Psi_{top}^{D+2} + c\dot{X}^{D+2} = \Psi^{D+2} \\
s\Psi^\mu &= 0 \\
s\Psi^{D+2} &= 0 \\
s\Psi^{D+1} &= \Phi \\
s\alpha &= -\dot{\Phi} \\
s\Phi &= 0
\end{aligned} \tag{3.5}$$

In agreement with the art of BRST invariant gauge-fixing, we introduce $D+2$ antighosts with ghost number (-1) and the associated Lagrange multipliers for the gauge conditions on the X^A 's. We also introduce an antighost $\overline{\Phi}$ with ghost number (-2) and its fermionic partner $\overline{\eta}$ with ghost number (-1) which we will use as a fermionic Lagrange multiplier for the gauge condition in the ghost sector. In this sector the action of s is

$$\begin{aligned}
s\overline{\Psi}^A &= b^A \\
sb^A &= 0 \\
s\overline{\Phi} &= \overline{\eta} \\
s\overline{\eta} &= 0
\end{aligned} \tag{3.6}$$

The gauge-fixing Lagrangian must be written as an s -exact term

$$\mathcal{L}^X + \mathcal{L}^{D+1} + \mathcal{L}^{D+2} + \mathcal{L}^\Phi = s\left(\overline{\Psi}^A(\dots)_A + \overline{\Phi}(\dots)\right) \tag{3.7}$$

For the gauge-fixing in the X -sector, we choose

$$\begin{aligned}
\mathcal{L}^X &= s \left(-\frac{e}{2} g_{\mu\nu} \bar{\Psi}^\nu b^\mu + g_{\mu\nu} \bar{\Psi}^\mu (\dot{X}^\nu + \bar{\eta} \Psi^\nu + \frac{e}{2} \Gamma_{\rho\sigma}^\nu \bar{\Psi}^\rho \Psi^\sigma) \right) \\
&= -\frac{e}{2} g_{\mu\nu} b^\mu b^\nu + g_{\mu\nu} b^\mu (\dot{X}^\nu + e \Gamma_{\rho\sigma}^\nu \bar{\Psi}^\rho \Psi^\sigma + \bar{\Psi}^\nu \eta + \bar{\eta} \Psi^\nu) \\
&\quad - \bar{\Psi}^\nu (g_{\mu\nu} \dot{\Psi}^\mu + e \partial_\rho g_{\mu\nu} \dot{X}^\nu \Psi^\rho) - \frac{e}{2} \partial_\nu \Gamma_{\mu\rho\sigma} \bar{\Psi}^\mu \Psi^\nu \bar{\Psi}^\rho \Psi^\sigma - \Gamma_{\mu\rho\sigma} \bar{\Psi}^\mu \eta \bar{\Psi}^\rho \Psi^\sigma
\end{aligned} \tag{3.8}$$

For the gauge-fixing in the e -sector, we choose

$$\mathcal{L}^{D+1} = -s \left(\bar{\Psi}^{D+1} e \left(m + \frac{b^{D+1}}{2} \right) \right) = -e \frac{(b^{D+1})^2}{2} + b^{D+1} (-me + \bar{\Psi}^{D+1} \eta) + 2m \bar{\Psi}^{D+1} \eta \tag{3.9}$$

After elimination of the field b^{D+1} , we obtain

$$\mathcal{L}^{D+1} \sim \frac{em^2}{2} + m \bar{\Psi}^{D+1} \eta \tag{3.10}$$

For the gauge-fixing in the X^{D+2} -sector, we choose

$$\begin{aligned}
\mathcal{L}^{D+2} &= s \left(\bar{\Psi}^{D+2} \left(-\frac{a}{2} b^{D+2} + X^{D+2} - C - \frac{1}{a} \frac{\bar{\Psi}^{D+1} \dot{\Psi}^{D+1}}{X^{D+2} - C} \right) \right) \\
&= -\frac{a}{2} (b^{D+2})^2 + b^{D+2} \left(X^{D+2} - C - a \frac{\bar{\Psi}^{D+1} \dot{\Psi}^{D+1}}{X^{D+2} - C} \right) \\
&\quad - \bar{\Psi}^{D+2} \left(\Psi^{D+2} - as \left(\frac{\bar{\Psi}^{D+1} \dot{\Psi}^{D+1}}{X^{D+2} - C} \right) \right)
\end{aligned} \tag{3.11}$$

a and C are arbitrarily chosen real numbers. After elimination of the field b^{D+2} , we find

$$\mathcal{L}^{D+2} \sim -\bar{\Psi}^{D+1} \dot{\Psi}^{D+1} + \frac{1}{2a} (X^{D+2} - C)^2 - \bar{\Psi}^{D+2} \left(\Psi^{D+2} - as \left(\frac{\bar{\Psi}^{D+1} \dot{\Psi}^{D+1}}{X^{D+2} - C} \right) \right) \tag{3.12}$$

The variable X^{D+2} can be eliminated by its algebraic equation of motion as well as the corresponding ghosts Ψ^{D+2} and $\bar{\Psi}^{D+2}$, after some field redefinitions. X^{D+2} is concentrated in a Gaussian way around the arbitrary scale C , with an arbitrary width a . We

are thus left with the propagating term for Ψ^{D+1} and $\bar{\Psi}^{D+1}$ which was missing in $\mathcal{L}^X$ and $\mathcal{L}^{D+1}$

$$\mathcal{L}^{D+2} \sim -\bar{\Psi}^{D+1} \dot{\Psi}^{D+1} \quad (3.13)$$

We finally choose the gauge-fixing in the ghost sector. To recover the full Lagrangian (3.2) and eventually identify the coordinate e as the einbein of the projection of the particle trajectory in the D -dimensional physical space-time, we need a term linear in $\bar{\eta}$ as well as another term to get rid of unwanted higher order fermionic terms. We define

$$\begin{aligned} \mathcal{L}^\Phi &= s(\bar{\Phi}(m\Psi^{D+1} - \Gamma_{\nu\rho\sigma}\bar{\Psi}^\rho\Psi^\sigma\Psi^\nu)) \\ &= \bar{\eta}(m\Psi^{D+1} - \Gamma_{\nu\rho\sigma}\bar{\Psi}^\rho\Psi^\sigma\Psi^\nu) + \bar{\Phi}(m\Phi - s(\Gamma_{\nu\rho\sigma}\bar{\Psi}^\rho\Psi^\sigma\Psi^\nu)) \end{aligned} \quad (3.14)$$

The dependence on the ghosts of ghosts Φ and $\bar{\Phi}$ is trivial: these fields decouple after a Gaussian integration. One has thus

$$\mathcal{L}^\Phi \sim m\bar{\eta}\Psi^{D+1} - \Gamma_{\nu\rho\sigma}\bar{\Psi}^\rho\Psi^\sigma\bar{\eta}\Psi^\nu \quad (3.15)$$

Adding all terms (3.8), (3.10), (3.13) and (3.15), we finally recognize that $\mathcal{L}^X + \mathcal{L}^{D+1} + \mathcal{L}^{D+2} + \mathcal{L}^\Phi$ is equivalent to the Lagrangian (3.2), modulo the elimination of auxiliary fields and the change of notation $(\eta, \bar{\eta}) \rightarrow (\alpha, \bar{\alpha})$. We have therefore shown the announced result: the $N = 2$ local supersymmetry of the Lagrangian describing spin-one particles is a residual symmetry coming from a topological model after a suitable gauge-fixing.

4 Derivation of Proca's equations from the (N=2)-supersymmetric action.

To verify the physical content of the model of the last section, we consider a flat space-time, and choose the gauge where the einbein and gravitini are constants over which we integrate. As explained in the first section, the Hamiltonian is

$$H = \frac{e_0}{2}(p^2 - m^2) + \bar{\alpha}_0 Q + \alpha_0 \bar{Q} \quad (4.1)$$

with

$$\begin{aligned} Q &= p_\mu \Psi^\mu + m \Psi^{D+1} \\ \overline{Q} &= p_\mu \overline{\Psi}^\mu + m \overline{\Psi}^{D+1} \end{aligned} \quad (4.2)$$

The matrices Ψ and $\overline{\Psi}$ satisfy the Clifford algebra

$$\{\Psi^A, \overline{\Psi}^B\}_+ = \eta^{AB} \quad , \quad \{\Psi^A, \Psi^B\}_+ = \{\overline{\Psi}^A, \overline{\Psi}^B\}_+ = 0 \quad (4.3)$$

for $A, B = 1, \dots, D+1$. Since the underlying gauge symmetry has Q and $\overline{Q}$ as generators, the physical states satisfy

$$Q|\phi\rangle = 0 \quad \overline{Q}|\phi\rangle = 0 \quad (4.4)$$

in addition to

$$(p^2 - m^2)|\phi\rangle = 0 \quad (4.5)$$

The Ψ and $\overline{\Psi}$ are generalizations of the Pauli matrices, and it is convenient to use a Schwinger type construction, in order to exploit directly their Clifford algebra structure. One introduces a spin vacuum $|0\rangle$ annihilated by the Ψ 's. Then, the $\overline{\Psi}$'s can be identified as their adjoints and act as creation operators. In the X representation, we can write a general state as

$$\begin{aligned} |\phi\rangle &= \left(\varphi_0 + \varphi_\mu \overline{\Psi}^\mu + \varphi_{\mu_1 \mu_2} \overline{\Psi}^{\mu_1} \overline{\Psi}^{\mu_2} + \dots + \varphi_{\mu_1 \dots \mu_D} \overline{\Psi}^{\mu_1} \dots \overline{\Psi}^{\mu_D} \right) |0\rangle \\ &+ \overline{\Psi}^{D+1} \left(\overline{\varphi}_0 + \overline{\varphi}_\mu \overline{\Psi}^\mu + \overline{\varphi}_{\mu_1 \mu_2} \overline{\Psi}^{\mu_1} \overline{\Psi}^{\mu_2} + \dots + \overline{\varphi}_{\mu_1 \dots \mu_D} \dots \overline{\Psi}^{\mu_D} \right) |0\rangle \end{aligned} \quad (4.6)$$

The wave functions $\varphi_{\mu_1 \dots \mu_p}(X)$ and $\overline{\varphi}_{\mu_1 \dots \mu_p}(X)$ are antisymmetric and it is useful to consider the differential forms

$$\begin{aligned} \varphi_p &= \frac{1}{p!} dX^{\mu_1} \dots dX^{\mu_p} \varphi_{\mu_1 \dots \mu_p}(X) \\ \overline{\varphi}_p &= \frac{1}{p!} dX^{\mu_1} \dots dX^{\mu_p} \overline{\varphi}_{\mu_1 \dots \mu_p}(X) \end{aligned} \quad (4.7)$$

for $0 \leq p \leq D$. The constraints (4.4) can be conveniently written as

$$d\varphi_p + im\overline{\varphi}_{p+1} = 0 \quad (4.8)$$

$$d^*\bar{\varphi}_p + im\varphi_{p-1} = 0 \quad (4.9)$$

$$d\bar{\varphi}_p = 0 \quad (4.10)$$

$$d^*\varphi_p = 0 \quad (4.11)$$

Where $d = dx^\mu \partial_\mu$ and d^* is its Hodge dual. One has also

$$(d^*d + dd^*)\varphi = -m^2\varphi \quad (d^*d + dd^*)\bar{\varphi} = -m^2\bar{\varphi} \quad (4.12)$$

These equations determine the independent degrees of freedom. When $m \neq 0$, they couple the two sectors of opposite chiralities. Moreover, when D is even, the first one contains $\frac{D}{2}$ forms, namely one scalar (φ_0), one vector (φ_1), ..., and one $(\frac{D}{2} - 1)$ -form ($\varphi_{\frac{D}{2}-1}$). The other one has a dual structure ($\bar{\varphi}_{\frac{D}{2}+1}, \dots, \bar{\varphi}_D$). For (φ_1), the constraints (4.4) can be rewritten:

$$\begin{aligned} \partial_\mu \bar{\varphi}^{\mu\nu} + im\varphi^\nu &= 0 \\ \partial_\mu \varphi^\mu &= 0 \\ \partial_{[\mu} \varphi_{\nu]} + im\bar{\varphi}_{\mu\nu} &= 0 \end{aligned} \quad (4.13)$$

Thus the vector wave function φ_1 satisfies Proca's equations, and describes a spin-one particle with mass m . It follows that the field equations of φ_1 and $\bar{\varphi}_1$ can be derived by minimizing Proca's Lagrangian

$$\begin{aligned} \mathcal{L}_{Proca} &= \frac{m}{2} \bar{\varphi}_{\mu\nu} \varphi^{\mu\nu} - \frac{i}{2} \varphi^{\mu\nu} (\partial_\mu \varphi_\nu - \partial_\nu \varphi_\mu) \\ &\quad - \frac{i}{2} \bar{\varphi}^{\mu\nu} (\partial_\mu \bar{\varphi}_\nu - \partial_\nu \bar{\varphi}_\mu) + m \bar{\varphi}_\mu \varphi^\mu \end{aligned} \quad (4.14)$$

When $m = 0$, the two sectors of opposite chiralities decouple. In each sector, the independent degrees of freedom are now one 0-form A_0 (with $\varphi_1 = dA_0$), one 1-form A_1 (with $\varphi_2 = dA_1$), ..., one $(D-2)$ -form A_{D-2} (with $\varphi_{D-1} = dA_{D-2}$). The φ_p 's are closed and co-closed, i.e. the A_p 's satisfy Maxwell's equations and are defined up to gauge transformations. Consequently, φ_2 can be identified with the field strength of a photon.

If we consider the case $D = 4$ and $m \neq 0$, the spectrum reduces to two scalars and two massive spin-one particles, and contains $8=2(1+3)$ degrees of freedom. For $m = 0$, we have two massless scalars and two massless vectors, so that we still have $8=2(1+1+2)$ independent degrees of freedom.

As an application of this formalism, we study the classical behavior of spinning particles in a curved space-time. We are interested in the approximation where the trajectory of the particle is classical, while the spin effects are visible as it would be the case in a Stern-Gerlach experiment. This situation occurs if the order of magnitude of the interaction energy between the spin and the curvature, which is essentially proportional to the space-time curvature times $\hbar$ (analogously to the interaction between the magnetic field and a magnetic moment due to the spin), is comparable to the kinematical energy of the particle. One must also measure the position of the particle on a domain much larger than its Compton wavelength. In this limit the position X^μ and momentum P_μ are ordinary numbers and the quantum Hamiltonian becomes simply a matrix built from the Ψ 's and $\bar{\Psi}$'s acting in the spin-space with coefficients depending on the classical position X and momentum P . The τ -dependence of the classical dynamics of the particle can be expressed by applying Hamilton-Jacobi's method with this matricial Hamiltonian. The only quantum effects are due to the spin interaction with the space-time curvature. (In a fully classical approximation, $\hbar = 0$, and the spin effects disappear, since all the fermionic operators are proportional to $\sqrt{\hbar}$.) One can always find a basis for the spin states, which depends on the space-time position and such that the Hamiltonian is diagonal. In this basis the spin value is conserved through evolution, i.e. the spin observables are parallelly transported along the trajectory. By diagonalization in spin space, H determines independent Hamilton-Jacobi's equations for each spin degree of freedom of the particle. For the spin-one case, we expect three different trajectories corresponding to the values 1, 0 and -1 for the projection of the spin on a spatial axis in the rest frame of the particle.

We consider the case of a Schwarzschild gravitational field in four dimensional space-time ($ds^2 = (1 - \frac{r_0}{r})dt^2 - (1 - \frac{r_0}{r})^{-1}dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2)$ with $r_0 = 2GM/c^2$.) We will compute the correction, due to the spin, to Einstein's formula predicting the

shift of the perihelion of a spinless point particle. For the other classical test of general relativity, i.e. the bending of light rays in a gravitational field, we will find that the wave vector of a polarized photon deviates from geodesic motions by a relative shift proportional to $\hbar$. These results are in agreement with the fact that a particle with an angular momentum interacts with the space-time curvature, as first pointed out by Papapetrou for a rotating body [16]. The advantage of a supersymmetric Hamiltonian is that it defines unambiguously the spin effects. Since we work to first non-trivial order in $\hbar$, we restore from now on the $\hbar$ dependence in the formulae. The matricial Hamilton-Jacobi's equation is obtained by replacing in the supersymmetric Hamiltonian the classical momentum p_μ by $\frac{\delta S}{\delta X^\mu}$ where $S[X^\mu, \tau]$ is the action of the classical trajectory of the particle in a given spin state, with arbitrarily chosen initial and final boundary conditions. Notice that keeping the lowest order in $\hbar$ means that we only retain the covariant derivative of the fermionic variables and not the curvature term. This yields

$$g^{\mu\nu} \frac{\delta S}{\delta X^\mu} \frac{\delta S}{\delta X^\nu} - m^2 + 2\hbar \frac{\delta S}{\delta X^\mu} \omega_{ab}^\mu \Sigma^{ab} + O(\hbar^2) = 0 \quad (4.15)$$

The space-time spin-connection ω is related to the space-time vierbein E and to Christoffel's symbol Γ

$$\omega_{\mu ab} = E_a^\alpha E_b^\beta \Gamma_{\alpha\mu\beta} \quad (4.16)$$

$$E_\alpha^a E_\beta^b \eta_{ab} = g_{\alpha\beta} \quad (4.17)$$

$$\Gamma_{\alpha\mu\beta} = \frac{1}{2}(\partial_\mu g_{\alpha\beta} + \partial_\beta g_{\alpha\mu} - \partial_\alpha g_{\beta\mu}) \quad (4.18)$$

The $\Sigma^{ab} = \frac{i}{2}(\bar{\Psi}^a \Psi^b - \bar{\Psi}^b \Psi^a)$ are the generators of the (reducible) 32-dimensional representation of the Lorentz group defined by the algebra (4.3) and acting on the states solving (4.6). If the matrix form of $\bar{\Psi}^5$ is chosen diagonal, the spin operators Σ^{ab} become block-diagonal with two independent sectors of opposite chiralities, corresponding to the eigenvalues 0 and 1 of $\bar{\Psi}^5 \Psi^5$, so the 32-dimensional representation splits into two independent 16-dimensional representation, each one containing five sectors of dimensions 1,4,6,4,1 corresponding respectively to 0-forms, 1-forms, 2-forms, 3-forms, and 4-forms. As explained above, the constraints imply that only two block-sectors made of one 0-form

and one 1-form sectors are independent wave-functions. The one-form sector, and the corresponding 4×4 Hamiltonian matrix, determine the dynamics of spin-one particles. Moreover, in a Schwarzschild metric with characteristic radius r_0 , the motion is planar, so one can separate the variables and write

$$S = -Et + L\varphi + S_r(r) \quad (4.19)$$

The spin-dependent part of Hamilton-Jacobi's equation is obtained by the substitution

$$p_\mu \omega_{ab}^\mu \Sigma^{ab} = \frac{r_0 E}{r^2} \Sigma^{01} - \frac{2L}{r^2} \left(1 - \frac{r_0}{r}\right)^{1/2} \Sigma^{13} \quad (4.20)$$

where

$$\Sigma^{01} = \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \Sigma^{13} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix} \quad (4.21)$$

By inserting (4.19) and (4.20) into Hamilton-Jacobi's equation (4.15), one obtains a matricial equation for $\frac{\partial S}{\partial r}$. The diagonalization can be done easily, and one gets three possibilities S_ϵ for the classical action, indexed by $\epsilon = 0, \pm 1$

$$\begin{aligned} \frac{E^2}{c^2} \left(1 - \frac{r_0}{r}\right)^{-1} - \left(\frac{L^2}{r^2} + m^2\right) - \left(1 - \frac{r_0}{r}\right) \left(\frac{\partial S_\epsilon}{\partial r}\right) \\ + 2\epsilon \frac{\hbar}{r^2} \left(L^2 \left(1 - \frac{r_0}{r}\right) - \left(\frac{r_0 E}{2c}\right)^2\right)^{1/2} = 0 \end{aligned} \quad (4.22)$$

(We have restored the dependence in the speed of light c .) The energy E and the angular momentum L are constants of motion of the particle. The values $\epsilon = 0, \pm 1$ correspond to the three possible projections of the spin along a given spatial axis in the rest frame of the particle. The case $\epsilon = 0$ corresponds to the geodesic trajectory followed by the scalar particle. Far from the Schwarzschild horizon, we can use the standard techniques of integration of Hamilton-Jacobi's equation to determine the three possibilities for the shift of the perihelion over a quasi-periodic trajectory. This amounts to replace L in the classical formulas [17] by an effective angular momentum L_ϵ defined by

$$L_\epsilon^2 = L^2 + 2\epsilon \hbar L \sqrt{1 - \left(\frac{r_0 E}{2Lc}\right)^2} \quad (4.23)$$

(Notice that near the horizon, unitarity breaks down). In the case of a massive particle, the shift of the perihelion is thus given by:

$$\delta\phi_\epsilon = \frac{3\pi}{2} \left(\frac{mcr_0}{L_\epsilon} \right)^2 \sim \delta\phi_0 \left(1 - \epsilon \frac{\hbar}{L} \sqrt{1 - \left(\frac{r_0 E}{2Lc} \right)^2} \right) \quad (4.24)$$

For a non-relativistic Z^0 orbiting quasi-tangentially to the sun at a speed of $10^5 m/s$, which is approximately the circular velocity around the sun, we find $|\delta\phi_+ - \delta\phi_0|/\delta\phi_0 \sim \hbar/L \sim 10^{-21}$, which is much too small to be detected.

The solutions of Hamilton-Jacobi's equation are continuous when $m \rightarrow 0$. However, in this limit the interpretation of its solution S is different. The particle is a photon following the laws of the geometrical optics, S is the eikonal of the light ray, and $\frac{\delta S}{\delta X^\mu}$ is its wave-vector. The solution $\epsilon = 0$ must then be rejected. In this massless case, one finds for the deflections of the two helicities $\epsilon = \pm 1$ the following formula

$$\delta\phi_\epsilon = \frac{2r_0\omega}{cL_\epsilon} \sim \delta\phi_0 \left(1 - \epsilon \frac{\hbar}{2L} \sqrt{1 - \left(\frac{r_0 E}{2Lc} \right)^2} \right) \quad (4.25)$$

where $\omega = \frac{E}{\hbar}$. For an optical photon of wavelength $\lambda = 7 \times 10^{-7} m$ (red) skimming past the sun, we find $|\delta\phi_+ - \delta\phi_0|/\delta\phi_0 \sim \frac{\hbar}{2L} = \frac{\lambda}{2R_{sun}} \sim 10^{-15}$. (Note that this ratio does not depend on $\hbar$: the gravitational field interacts classically with the two polarizations of the electromagnetic field.) However, this doubling of Einstein's rings is too small to be detected.

References

- [1] F.A. Berezin and M.S. Marinov, *JETP Lett.* **21** (1975) 320 and *Ann. Phys. NY* **104** (1977) 336; R. Casalbuoni, *Nuovo Cimento* **33A** (1976) 389 and *Phys. Lett.* **62B** (1976) 49; A. Barducci, R. Casalbuoni and L. Lusanna, *Nuovo Cimento* **35A** (1976) 377; L. Brink, S. Deser, B. Zumino, P. di Vecchia and P.S. Howe *Phys. Lett.* **64B** (1976) 43.
- [2] E. Witten, *Nucl. Phys.* **B202** (1982) 253; L. Alvarez-Gaumé, *Comm. Math. Phys.* **90** (1983) 161; D. Friedan and P. Windey, *Nucl. Phys.* **B235** (1984) 395;
- [3] L. Brink and J.H. Schwarz, *Nucl. Phys.* **B121** (1977) 285; L. Brink, P. di Vecchia and P.S. Howe, *Nucl. Phys.* **B118** (1977) 76 and *Phys. Lett.* **65B** (1976) 471; S. Deser and B. Zumino *Phys. Lett.* **65B** (1976) 369.
- [4] A.M. Polyakov, *Phys. Lett.* **103B** (1981) 211.
- [5] P.A.M. Dirac. Lectures on Quantum Mechanics, (Belfer Graduate School of Science, Yeshiva University; New York; 1964); M. Henneaux and C. Teitelboim, Quantization of Gauge Systems (Princeton University Press; 1992).
- [6] W. Siegel, Introduction to String Field Theory. (World Scientific; 1988).
- [7] A.M. Polyakov, Gauge Fields and Strings (Harwood Academic Publishers; 1987); Vl. S. Dotsenko, *Nucl. Phys.* **B285** (1987) 45.
- [8] R.H. Rietdijk and J.W. van Holten, *Class. Quantum Grav.* **7** (1990) 247.
- [9] D. Birmingham, M. Blau, M. Rakowski and G. Thomson, *Physics Reports* **209** (1991) 129.
- [10] L. Baulieu and I.M. Singer, *Comm. Math. Phys.* **125** (1989) 227.
- [11] L. Baulieu and I.M. Singer, *Nucl. Phys. Proc. Suppl.* **5B** (1988) 12.
- [12] P. Howe, S. Penati, M. Pernici and P. Townsend, *Phys. Lett.* **215B** (1988) 555.
- [13] R. Marnelius and U. Martensson, *Nucl. Phys.* **B335** (1990) 395; U. Martensson, Preprint Goteborg-92-3 (Jan. 92).
- [14] P. Van Nieuwenhuizen, *Phys. Rep.* **68C** (1981) 189.

- [15] R.H. Rietdijk and J.W. van Holten, *Class. Quantum Grav.* **10** (1993) 575.
- [16] A. Papapetrou, *Proc. Roy. Soc. London* **A209** (1951) 248.
- [17] L.D. Landau, *The Classical Theory of Fields*. (Pergamon Press; London; 1971).